

Preen

LAWN

FALL WEED CONTROL
PLUS FERTILIZER



COVERS 5,000 SQ. FT.



Preen

LAWN

FALL WEED CONTROL PLUS FERTILIZER

- Kills dandelions, chickweed, clover (over 200 weeds)
- Strengthens roots for winter hardiness
- Apply in fall to reduce spring weeds



SPECIMEN LABEL

COVERS 5,000 SQ. FT.



LAWN
FALL WEED CONTROL
PLUS FERTILIZER

Preen

COVERS
5,000 SQ. FT.
Net Wt. 16 lbs. (7.26 kg)

Keep out of reach of children

CAUTION

See back panel for additional Precautionary Statements.

26-2-12



ACTIVE INGREDIENTS:	
2,4-D	0.76%
Mecoprop-p (MCP-p)	0.25%
Dicamba	0.07%
OTHER INGREDIENTS:	98.92%
TOTAL	100.00%

THIS PRODUCT CONTAINS:
15.2 lbs. 2,4-dichlorophenoxyacetic acid equivalent per ton.
3.4 lbs. (1+1R)-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per ton.
1.4 lbs. 3,5-dichloro-*m*-cresamic acid equivalent per ton.

Former Specific by ADAC Methods
Contains single isomer form of Mecoprop-p.

- Kills dandelions, chickweed, clover (over 200 weeds)
- Strengthens roots for winter hardiness
- Apply in fall to reduce spring weeds



COVERS
5,000 SQ. FT.
 Net Wt. 16 lbs. (7.26 kg)

Keep out of reach of children

CAUTION

See back panel for additional Precautionary Statements.

26-2-12



ACTIVE INGREDIENTS:	
2,4-D.....	0.76%
Mecoprop-p (MCPP-p).....	0.17%
Dicamba.....	0.07%
OTHER INGREDIENTS:	99.00%
TOTAL	100.00%

THIS PRODUCT CONTAINS:
 15.2 lbs. 2,4-dichlorophenoxyacetic acid equivalent per ton.
 3.4 lbs. (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per ton.
 1.4 lbs. 3,6-dichloro-o-anisic acid equivalent per ton.

Isomer Specific by AOAC Methods
 Contains single isomer form of Mecoprop-p.

SPECIMEN LABEL

Preen®

LAWN

FALL WEED CONTROL PLUS FERTILIZER

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until spray has dried or dust has settled.

Use Precautions

Do not use on dichondra, carpetgrass or St. Augustinegrass.

Do not use this product where desirable clovers are present or allow the material to drift onto flowers, vegetables or ornamental shrubbery. Do not apply in areas underlaid by roots of desirable trees or shrubs. Apply when the air is calm. Do not apply to newly seeded lawns. During application, close spreader openings when turning.

Instructions

Preen Lawn Fall Weed Control Plus Fertilizer is a Trimec® complex, a combination of three proven weed killers — 2,4-D, Mecoprop-p and Dicamba. Together, they control a wide range of lawn weeds, including dandelion, chickweed, knotweed, plantain, henbit, spurge and many others commonly found in home lawns. In addition, a 26-2-12 fertilizer provides a supply of plant nutrients. Possible retreatment may be necessary if (a) lawn is heavily overrun by weeds, or, (b) if adverse low moisture conditions prevail and weeds are in a state of poor growth.

Apply 3.2 lbs. of product per 1,000 sq. ft. of turfgrass. A second application may be made in approximately 30 days.

The maximum application rate to turf is 1.2 lbs. 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

When to Apply

Apply when broadleaf weeds are actively growing in spring or early fall.

What to Expect

Weed stems and leaves will begin to twist and curl in a few days, but a month or more may be required for complete control. This product will not prevent weed seeds from producing new plants.

1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least 1 to 2 days before application to sustain moisture until the next watering (see step 5).
3. Apply when weeds are young and actively growing, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to hold the material and prevent dust drift onto nontarget plants.
4. Avoid applying when temperatures are consistently above 90°F.
5. Do not wash from weed leaves for 1 to 2 days after application. At this time, a thorough watering should be made.

Broadleaf Weeds

Annual yellow sweetclover	Cocklebur	Orange hawkweed	Spotted spurge
Aster	Common mullein	Oriental cocklebur	Spurweed
Austrian fieldcress	Creeping jenny	Oxalis	Stinging nettle
Bedstraw	Cudweed	Parsley-piert	Strawberry, India mock
Beggarticks	Curly dock	Parsnip	Tall nettle
Betony, Florida	Daisy: English, oxeye	Pearlwort	Tall vervain
Bindweed, field	Daisy fleabane	Pennycress	Tansy ragwort
Bird vetch	Dandelion	Pepperweed	Tansy mustard
Bitter wintercress	Dichondra	Pigweed	Tanweed
Bittercress, hairy	Dogfennel	Pineywoods bedstraw	Thistle
Bitterweed	Dollarweed	Plains coreopsis	Trailing crownvetch
Black-eyed Susan	False dandelion	Plantain	Tumble mustard
Black medic	Falseflax	Poison ivy	Tumble pigweed
Black mustard	False sunflower	Poison oak	Velvetleaf
Blackseed plantain	Fiddleneck	Pokeweed	Venice mallow
Blessed thistle	Florida pusley	Poorjoe	Virginia buttonweed
Bloodflower	Frenchweed	Prairie sunflower	Virginia creeper
milkweed	Galinsoga	Prickly lettuce	Virginia pepperweed
Blue lettuce	Goldenrod	Prickly sida	Wavyleaf bullthistle
Blue vervain	Ground ivy	Prostrate knotweed	Western clematis
Bracted plantain	Gumweed	Prostrate pigweed	Western salsify
Bristly oxtongue	Hairy fleabane	Prostrate spurge	White mustard
Broadleaf dock	Hawkweed	Prostrate vervain	Wild aster
Broadleaf plantain	Healall	Puncturevine	Wild buckwheat
Broomweed	Heartleaf drymary	Purslane, common	Wild carrot
Buckhorn	Heath aster	Ragweed	Wild four-o'clock
Buckhorn plantain	Hedge bindweed	Red sorrel	Wild garlic
Bulbous buttercup	Hedge mustard	Redroot pigweed	Wild geranium
Bull thistle	Henbit	Redstem filaree	Wild lettuce
Bullnettle	Hoary cress	Rough cinquefoil	Wild marigold
Burclover	Hoary plantain	Rough fleabane	Wild mustard
Burdock	Hoary vervain	Russian pigweed	Wild onion
Burning nettle	Horsenettle	Russian thistle	Wild parsnip
Burweed	Jimsonweed	Scarlet pimpernel	Wild radish
Buttercup	Knawel	Scotch thistle	Wild rape
Buttonweed	Knotweed	Sheep sorrel	Wild strawberry
Canada thistle	Kochia	Shepherdspurse	Wild sweet potato
	Lambsquarters		

SPECIMEN LABEL

Calibration Instructions

To provide proper distribution, calibrate spreader before application. For 3.2 lbs. of product per 1,000 sq. ft. of turfgrass: Check your spreader on 310 sq. ft. and adjust to apply at the rate of 1 lb. of product per 310 sq. ft.


Distribution and Spreader Patterns

Always apply with a properly calibrated rotary or drop type fertilizer spreader to distribute the granules evenly. Never apply by hand.

Coverage - To determine how much you need to apply, multiply the length of your lawn by its width. This equals your total square footage. Deduct all non-lawn areas (house, gardens, etc.). One bag covers up to 5,000 sq. ft.

Note: Supplemental applications of lawn fertilizer are needed to achieve a minimum 1 lb. nitrogen per 1,000 sq. ft.

Suggested Spreader Settings




Apply in a single pass at a rate of 3.2 lbs. per 1,000 sq. ft. Start on the outside border and apply in a circular path working toward the center. Overlap the coverage slightly, and shut off spreader before making hard turns.

Broadcast Spreaders

Setting

Agri-Fab® Rotary	4
Cyclone® Rotary	4
Earthway® Rotary	14
Greenview® Jet-Spred® SS1	14
Lesco® Rotary (dial A-S)/(number dial)	H /16
Red Devil® Rotary (1-20)	5
Republic® Rotary	4
Scotts® EasyGreen®	27
Scotts® EdgeGuard®	4 1/2
Scotts® SpeedyGreen® 2000, 3000	4 1/2
Sears® Craftsman Rotary	5
Sta-Green® Rotary (1-10)	4
True Temper® Rotary	4
Vigoro® Rotary (1-10)	4



Apply in a single pass at a rate of 3.2 lbs. per 1,000 sq. ft. Start on the outside border and apply in a rectangular path working back and forth across the length of the lawn. Overlap the coverage slightly, and shut off spreader before making hard turns.

Drop Spreaders

Setting

Agri-Fab® Drop	5
Greenview® Drop-Spred® XP1	18
Red Devil® Drop (1-5)	4 1/2
Republic® Drop	6
Scotts® Drop AccuGreen® (2-18)	5
Scotts® Drop PF1 (2-18)	5
Sta-Green® Drop (1-5)	4 1/2



If your spreader model does not appear above, go to our website at www.preen.com, or call toll-free (800) 233-1067, 8 A.M. - 5 P.M. EDT, Monday - Friday, for additional model settings.

Carolina geranium	Lespedeza	Slender plantain	Wild vetch
Carpetweed	Mallow	Smallflower	Woodsorrel
Catchweed bedstraw	Matchweed	galinsoga	Woolly croton
Catnip	Mexicanweed	Smooth dock	Woolly morningglory
Catsear	Milk vetch	Smooth pigweed	Woolly plantain
Chickweed: common,	Morningglory	Sorrel	Wormseed
mouseear	Mouseear hawkweed	Sowthistle	Yarrow
Chicory	Mugwort	Spanishneedles	Yellow rocket
Cinquefoil	Musk thistle	Speedwell	Yellowflower
Clover: crimson,	Mustard	Spiny amaranth	pepperweed
hop, red, strawberry,	Narrowleaf plantain	Spiny cocklebur	& other broadleaf
sweet, white	Narrowleaf vetch	Spiny sowthistle	weeds
Cockle	Nettle	Spotted catsear	

Guaranteed Fertilizer Analysis:

Total Nitrogen (N)	26%
2.0% Ammoniacal Nitrogen	
5.2% Water Insoluble Nitrogen	
13.6% Urea Nitrogen	
5.2% Other Water Soluble Nitrogen*	
Available Phosphate (P ₂ O ₅)	2%
Soluble Potash (K ₂ O)	12%
Total Sulfur (S)	4.0%
4.0% Combined Sulfur (S)	
Nutrient Sources: Ammonium Phosphate, Ammonium Sulfate, Methylene Ureas, Urea, Muriate of Potash, Sulfate of Potash.	
Chlorine (Cl) not more than	10.0%
* 5.2% Slowly Available Nitrogen from Methylene Ureas.	F699

Warranty

The distributor warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to distributor, and buyer assumes the risk of any such use.

PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals

CAUTION Harmful if swallowed. Do not inhale dust. Do not allow contact with skin, eyes or clothing. When using this product, wear long-sleeved shirt, long pants, socks and shoes.

After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water.

FIRST AID

IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact (888) 208-1368 for emergency medical treatment information.

Environmental Hazards

Drift or runoff may adversely affect nontarget plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwater. Do not apply when weather conditions favor drift from target area.

Storage and Disposal

Storage: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. **Container Disposal: If empty:** Do not reuse this container. Place in trash or offer for recycling if available. **If partially filled:** Call your local solid waste agency or toll free number (1-800-CLEANUP) for disposal instructions. Never place unused product down any indoor or outdoor drain.

EPA Reg. No. 2217-660-961

(L) (D)

EPA Est. 961-PA-01:961-IL-01

Superscript used is first letter of lot number.

CAS Registry Numbers:

2,4-D CAS No. 94-75-7

Mecoprop-p CAS No. 16484-77-8

Dicamba CAS No. 1918-00-9



PROOF OF PURCHASE

PREEN GUARANTEE

Lebanon Seaboard Corporation guarantees you will be satisfied with the performance of this product when used as directed, or your money back. If not satisfied, describe the nature of your problem, and send with proof of purchase to:

LEBANON SEABOARD CORPORATION
1600 EAST CUMBERLAND STREET
LEBANON, PA 17042



1-800-233-1067



www.preen.com

Trimec® is a registered trademark of PBI/Gordon Corporation.

Preen® is a registered trademark of Lebanon Seaboard Corporation.

6/05

Information regarding the levels of metals in this product is available on the Internet at <http://www.regulatory-info-lebsea.com>

SPECIMEN LABEL