

APPLY
- IN -
SPRING

Preen[®] One

LAWN CARE

EASY ONE-STEP APPLICATION
FOR SPRING LAWN CARE!



STOPS
Crabgrass All Season.

Can be applied
up to 4 weeks after
crabgrass emerges.



KILLS
Dandelions And
Most Lawn Weeds*:

Kills 250 weeds*
to the root.



FEEDS
Your Lawn.

Feeds the lawn
for 8 weeks.

COVERS 2,500 SQ.FT. CUBRE 232.25 m² (2,500 pies²)

*SEE USE DIRECTIONS FOR COMPLETE LIST OF TREATED WEEDS.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See back panel for First Aid and additional Precautionary Statements.

**MANTENGA ALEJADO DEL ALCANCE DE LOS NIÑOS
PRECAUCIÓN**

Consulte el panel posterior para obtener información sobre primeros auxilios y otras declaraciones de precaución adicionales.

New York State: Do not exceed one application at the maximum rate of 3.6 pounds of product per 1,000 sq.ft. per year (0.25 lb. of dithiopyr per acre per year).

| | |
|--------------------------------|----------|
| ACTIVE INGREDIENTS: | |
| 2,4-D, 2-ethylhexyl ester..... | 1.108% |
| Mecoprop-p | 0.167% |
| Dicamba | 0.071% |
| Dithiopyr | 0.160% |
| OTHER INGREDIENTS:..... | 98.494% |
| TOTAL | 100.000% |

THIS PRODUCT CONTAINS:
14.70 lb 2,4-dichlorophenoxyacetic acid equivalent per ton or 0.735% (CAS 1928-43-4)
3.34 lb (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per ton or 0.167% (CAS 16484-77-8)
1.42 lb 3,6-dichloro-o-anisic acid equivalent per ton or 0.071% (CAS 1918-00-9)
3.20 lb dithiopyr per ton or 0.160% (CAS 97886-45-8)
Isomer Specific By AOAC Methods.

24-0-6

**NET WT. 9 LBS
PESO NETO (4,08 KG)**

EASY ONE-STEP APPLICATION FOR SPRING LAWN CARE!



STOPS
Crabgrass All Season.

Can be applied up to 4 weeks after crabgrass emerges.



KILLS
Dandelions And Most Lawn Weeds*:

Kills 250 weeds* to the root.



FEEDS
Your Lawn.

Feeds the lawn for 8 weeks.

COVERS 2,500 SQ.FT. CUBRE 232.25 m² (2,500 pies²)

*SEE USE DIRECTIONS FOR COMPLETE LIST OF TREATED WEEDS.

KEEP OUT OF REACH OF CHILDREN CAUTION

See back panel for First Aid and additional Precautionary Statements.

MANTENGA ALEJADO DEL ALCANCE DE LOS NIÑOS PRECAUCIÓN

Consulte el panel posterior para obtener información sobre primeros auxilios y otras declaraciones de precaución adicionales.

New York State: Do not exceed one application at the maximum rate of 3.6 pounds of product per 1,000 sq.ft. per year (0.25 lb. of dithiopyr per acre per year).

ACTIVE INGREDIENTS:

| | |
|--------------------------------|----------|
| 2,4-D, 2-ethylhexyl ester..... | 1.108% |
| Mecoprop-p | 0.167% |
| Dicamba | 0.071% |
| Dithiopyr | 0.160% |
| OTHER INGREDIENTS:..... | 98.494% |
| TOTAL | 100.000% |

THIS PRODUCT CONTAINS:

- 14.70 lb 2,4-dichlorophenoxyacetic acid equivalent per ton or 0.735% (CAS 1928-43-4)
 - 3.34 lb (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per ton or 0.167% (CAS 16484-77-8)
 - 1.42 lb 3,6-dichloro-o-anisic acid equivalent per ton or 0.071% (CAS 1918-00-9)
 - 3.20 lb dithiopyr per ton or 0.160% (CAS 97886-45-8)
- Isomer Specific By ADAC Methods.

24-0-6

**NET WT. 9 LBS
PESO NETO (4,08 KG)**

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear protective eyewear. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

| First Aid | |
|---|--|
| If in eyes: | <ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice. |
| If swallowed: | <ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person. |
| If on skin or clothing: | <ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice. |
| Have the product container or label with you when calling a poison control center or doctor or going for treatment. | |
| For additional information in case of emergency, call toll free (888)-208-1368 | |

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn will help to prevent run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.

USE INSTRUCTIONS

This product is formulated to provide control of most annual grasses and certain broadleaf weeds* commonly found in home lawns.

This product contains three components: a preemergent herbicide, a lawn fertilizer, and a post emergent broadleaf weed* killer.

- The preemergent herbicide can be applied to established turfgrass to control crabgrass, goosegrass, foxtails, and other undesirable annual grasses.
- The fertilizer contains nutrients for turfgrass uptake.
- The post emergent broadleaf herbicide will control over 250 lawn weeds* including dandelion, chickweed, thistle and clover.

A list of broadleaf weeds is presented on this label.

*See use directions for complete list of treated weeds.

Controls 253 weeds!

| | | | |
|-----------------------|------------------------|------------------------------------|----------------------------|
| Amaranth, purple | Charlie) | Mugwort | Speedwell, bi-lobed |
| Amaranth, slender | Cupid shaving Brush | Mustard, ball | Speedwell, common |
| Aster, heath | Cudweed | Mustard, black | Speedwell, corn |
| Aster, white heath | Daisy, English | Mustard, blue | Speedwell, ivyleaf |
| Aster, purple | Daisy, oxeye | Mustard, hedge | Speedwell, Persian |
| Aster, white prairie | Dandelion, common | Mustard, Indian | Speedwell, purslane |
| Artichoke | Dandelion, false | Mustard, tansy | Speedwell, slender |
| Austrian fieldcress | Day flower | Mustard, tumble | Speedwell, snow |
| Bedstraw, Catchweed | Deadnettle | Mustard, white | Speedwell, thymeleaf |
| Bedstraw, smooth | Dichondra | Mustard, wormseed | Spiny amaranth |
| Bedstraw, Northern | Dock, broadleaf | Nettle, stinging (burning) | Spurge, spotted |
| Bedstraw, pineywoods | Dock, curly | Nettle, tall | Spurge, prostrate |
| Beggarticks, hairy | Dogbane | Nettle, wood | Spurweed |
| Betony, Florida | Dogfennel | Parsley-piert | Star of Bethlehem |
| Bindweed, Field | Dollarweed (Pennywort) | Parsnip | Strawberry, India mock |
| Bindweed, Hedge | Dove weed | Pearwort | Tansy ragwort |
| Bitter Wintercress | Elderberry | Pennycress (frenchweed) | Tanweed |
| Bittercress, hairy | Falseflax | Pepperweed, perennial | Thistle, blessed |
| Bitterweed | False sunflower | Pepperweed, Virginia (Peppergrass) | Thistle, bull |
| Black-eyed Susan | Fiddleneck | Pigweed, prostate | Thistle, Canada |
| Black Medic | Flaree, redstem | Pigweed, redroot | Thistle, fiodman (prairie) |
| Blood flower milkweed | Flaree, whitestem | | Thistle, musk |

USE RESTRICTIONS

The use of Preen One Lawncare on lawns and ornamental turfgrasses that are not well established, or have been weakened by stress due to mechanical operations, chemical injury, disease, insects, pests, excessive temperatures, or lack of moisture may increase the chances for turfgrass injury. Allow turfgrass weakened by these stresses to recover before application of this product.

Do not use this product on or near desirable plants, including within the drip line of the roots of desirable trees and shrubs, since injury may result.

Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches.

Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans.

Do not apply when windy.

Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

Do not exceed specified dosage for any area.

Do not use clippings from treated areas for composting or for mulching vegetables or fruit trees.

WHEN TO APPLY

This product may be applied to established lawns as a single application or as split applications in the spring, early summer, or fall.

For the best results in the spring, apply this product one to two (1 to 2) weeks before the expected germination of crabgrass. When the period of germination for crabgrass and other weeds listed on this label extends into early summer, then a second application can be made at 6 to 10 weeks after the initial application.

For control of crabgrass and other listed grasses: This product is a preemergent herbicide to control crabgrass (smooth, southern and large) and other susceptible annual grasses. This product may also provide early post emergent control of crabgrass. However, post emergent crabgrass control will be obtained only when this product is applied before the first tiller stage of crabgrass.

For broadleaf weed control: Best results are obtained during the spring and fall when broadleaf weeds are present, young and actively growing under adequate soil moisture and favorable temperatures (50° to 80°F).

HOW TO APPLY

This product needs to stick to the leaves of broadleaf weeds and should be applied when the grass is wet from dew or rain; poor results will occur if it is applied to dry lawns. The lawn should not be watered or disturbed for 24 hours after application. Delay mowing for 3 days after application.

Apply this product when the wind is calm. Avoid spreading granules onto flowers, vegetables, and ornamental shrubbery or desirable plants.

HOW MUCH TO APPLY

Kentucky bluegrass, fescue spp., perennial ryegrass, common bermudagrass, and zoysiagrass. Apply 3.6 pounds of product per 1,000 square feet of established turfgrass.

Hybrid bermudagrass, bahiagrass and buffalograss. Apply 3 pounds of product per 1,000 square feet of established turfgrass.


New York State: Do not exceed one application at the maximum rate of 3.6 pounds of product per 1,000 sq. ft. per year (0.25 lb. active ingredient of dithiopyr per acre per year).

Other States: The maximum broadcast application rate is 3.6 pounds of product per 1,000 sq. ft. per application. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 7.2 pounds of product per 1,000 sq.ft.

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation, and evenness of terrain may require slightly different settings for desired coverage.

Broadcast Spreader Delivers 3.6 lbs./1,000 sq. ft.

| | |
|--|-------|
| Agri-Fab® Rotary | 4 |
| Cyclone® Rotary | 3 1/2 |
| Earthway® Rotary | 14 |
| Greenview® Jet-Spred® SS1 | 14 |
| Lesco® Rotary (dial A-S)/(number dial) | H/14 |
| Red Devil® Rotary (1-20) | 5 |
| Republic® Rotary | 4 |
| Scotts® EasyGreen® | 23 |
| Scotts® EdgeGuard® | 4 1/2 |
| Scotts® SpeedyGreen® 2000, 3000 | 4 1/2 |
| Sears® Craftsman Rotary | 4 |
| Sta-Green® Rotary (1-10) | 2 1/2 |
| True Temper® Rotary | 3 1/2 |
| Vigoro® Rotary (1-10) | 2 1/2 |



Broadcast Spreader: Apply in a single pass by starting on the outside border and apply in a circular path working toward the center.

| | | | |
|------------------------------------|--------------------------|---------------------------------|------------------------------|
| Blue flower milkweed | Platane, whitestem | Pigweed, Russian | Thistle, niobr |
| Blue lettuce | Fleabane, daisy (annual) | Pigweed, smooth | Thistle, plumeless |
| Brassbuttons | Fleabane, rough | Pigweed, tumble | Thistle, Russian |
| Bristly oxtongue | Galinsoga, hairy | Pineapple weed | Thistle, Scotch |
| Broomweed | Galinsoga, smallflower | Plains coreopsis (tickseed) | Thistle, wavyleaf |
| Buckhorn | Goldenrod | Plantain, blackseed | Thistle, yellow (horrible) |
| Bullnettle | Ground ivy | Plantain, bracted | Trailing crownvetch |
| Burclover | Groundsel | Plantain, broadleaf | Velvetleaf |
| Burdock | Gumweed | Plantain, buckhorn | Vervain, blue |
| Burweed, lawn (spurweed) | Hairy fleabane | Plantain, hoary | Vervain, hoary |
| Buttercup, bulbous | Hawkweed, meadow | Plantain, narrowleaf | Vervain, prostrate |
| Buttercup, creeping | Hawkweed, mouseear | Plantain, slender | Vervain, tall |
| Buttonweed, common | Hawkweed, orange | Plantain, woolly | Vetch, bird |
| (poorjoe) | Hawkweed, yellow | Poison ivy | Vetch, hairy |
| Buttonweed, Virginia | Healall | Poison oak | Vetch, milk |
| Carolina geranium (wild) | Heartleaf drymary | Pokeweed | Vetch, narrowleaf |
| Carpetweed | Hemp | Prairie Sunflower | Vetch, wild |
| Catnip | Henbit | Prickly lettuce (compass plant) | Virginia creeper |
| Catsear, spotted (false dandelion) | Hoary cress | Prickly sida | Western clematis |
| Chamber bitter | Horsenettle | Prostrate knotweed | Western salsify |
| Chickweed, common | Horseweed | Puncturevine | Wild aster |
| Chickweed, mouseear | Jimsonweed | Purslane, common | Wild buckwheat |
| Chickweed, sticky | Johnny-Jumpup violet | Pusley, Brazil | Wild carrot |
| Chicory | Knawel | Pusley, Florida | Wild four-o'clock |
| Cinquefoil | Knotweed, prostrate | Pusley, large flower | Wild garlic |
| Clover, crimson | Kochia | Ragweed, bur | Wild lettuce |
| Clover, alyce | Lambsquarters | Ragweed, common | Wild marigold |
| Clover, hop | Lespedeza | Ragweed, lanleaf | Wild mustard (yellow rocket) |
| Clover, rabbit foot | Mallow, alkali | Ragweed, western | Wild onion |
| Clover, red | Mallow, bristly | Redstem filaree | Wild parsnip |
| Clover, strawberry | Mallow, common | Rough cinquefoil | Wild radish |
| Clover, yellow sweet | Mallow, dwarf | Scarlet pimpernel | Wild rape |
| Clover, white sweet | Mallow, venice | Shepardspurse | Wild strawberry |
| Clover, white | Marcela | Smartweed, ladysthumb | Wild sweet potato |
| Cockle | Matchweed | Smartweed, pale | Wild violet |
| Cocklebur | Mexicanweed | Smartweed, Pennsylvania | Woodsorrel, creeping |
| Cocklebur, oriental | Moneywort | Smooth chaff-flower | Woodsorrel, violet |
| Cocklebur, spiny | Morningglory, bigroot | Smoothdock | Woodsorrel, yellow (Oxalis) |
| Common mullein | Morningglory, ivyleaf | Sorrel, red (sheep) | Woolly croton |
| Corn Spurry | Morningglory, Mexican | Sowthistle, annual | Wormseed |
| Creeping Beggarweed | Morningglory, red | Sowthistle, spiny | Yarrow, common |
| Creeping Jenny (creeping) | Morningglory, tall | Spanish needles | Yarrow, western |
| | Morningglory, woolly | | Yarrow, yellow |

WHERE TO USE
Preen One LawnCare may be used on the following established turf species:
Established Cool Season Turfgrass: Kentucky bluegrass, perennial ryegrass, fine fescues, tall fescue
Established Warm Season Turfgrass: bahiagrass, bermudagrass, buffalograss, zoysiagrass

Do not use this product on dichondra, carpetgrass or colonial bentgrass.
Do not use this product on St. Augustinegrass in Florida.
Do not use this product on turfgrass where desirable clovers are present.

If the lawn has been seeded recently, wait until after the third mowing before using this product. Do not seed, overseed or sprig the treated area for 10 weeks after application.

EPA Reg. No. 961-419


(L) (D)

EPA Est. No. 961-PA-01:961-IL-01
Superscript used is first letter of lot number.

Manufactured by Lebanon Seaboard Corporation
1600 East Cumberland Street | Lebanon, PA 17042

Preen® is a registered trademark
of Lebanon Seaboard Corporation.

Information regarding the contents and levels of metals in this product is available on the Internet at:
<http://www.regulatory-info-lebsea.com>



Drop Spreader: First make two passes along the top of the area you'll be covering. Then make two passes along the bottom area. Then move back and forth along the long part of the lawn. On each pass, as you reach the strips at the top and bottom of the area to be covered, shut off the spreader. It's also important to shut off the spreader when you are stopped.

Drop Spreader Delivers 3.6 lbs./1,000 sq. ft.

| | |
|--------------------------------|-------|
| Agri-Fab® Drop | 4 |
| Greenview® Drop-Spred® XP1 | 16 |
| Red Devil® Drop (1-5) | 3 1/2 |
| Republic® Drop | 5 |
| Scotts® Drop AccuGreen® (2-18) | 6 |
| Scotts® Drop PF1 (2-18) | 6 |
| Sta-Green® Drop (1-5) | 3 1/2 |

Note: if spreader model does not appear above please call toll free (800)233-1067, 8 am – 5 pm ET, Monday – Friday for additional model settings.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.
PESTICIDE STORAGE: Keep in original container in a cool, dry area, away from extreme heat and direct sunlight. Do not store near food feed, seeds or other pesticides.
PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

WARRANTY

NOTE To the extent consistent with applicable law, Buyer assumes all responsibility for safety and performance if this product is not used according to the directions.


NOTE: Supplemental applications of lawn fertilizer are needed to achieve a minimum 1 lb. nitrogen per 1,000 sq. ft.

Guaranteed Analysis:

| | |
|--|------|
| Total Nitrogen (N) | 24% |
| 2.8% Ammoniacal Nitrogen | |
| 1.8% Water Insoluble Nitrogen* | |
| 16.4% Urea Nitrogen | |
| 3.0% Other Water Soluble Nitrogen* | |
| Soluble Potash (K ₂ O)..... | 6% |
| Sulfur (S) | 3.3% |
| 3.3% Combined Sulfur (S) | |

Derived from: Ammonium Sulfate, Methylene Ureas, Urea, Muriate of Potash.

* 4.8% Slowly Available Nitrogen from Methylene Ureas. F699



PREEN GUARANTEE

To the extent consistent with applicable law, Lebanon Seaboard Corporation guarantees you will be satisfied with the performance of this product when used as directed, or your money back. If not satisfied, describe the nature of your problem, and send with proof of purchase to:

LEBANON SEABOARD CORPORATION
1600 EAST CUMBERLAND STREET, LEBANON, PA 17042

1-800-233-1067 **www.preen.com**